DOCKET NO.: AM100299 CON US/WYNC-0677

**Application No.:** 10/663,533

Office Action Dated: April 14, 2004

This listing of claims will replace all prior versions, and listings, of claims in the application.

**PATENT** 

## Listing of Claims

Claims 1 to 25 (cancelled)

26. (previously presented) A method of treating a subject suffering from a condition selected from the group consisting of neurodegenerative disease, eating disorders, disorders of thermoregulation, sleep dysfunction and sexual dysfunction, comprising the step of:

providing to the subject suffering from said condition, a therapeutically effective amount of a compound of formula I

$$R^{1}O$$
 $R^{1}O$ 
 $R^{2}$ 
 $R^{2}$ 

wherein

R<sup>1</sup> is a straight-chained alkyl of 1 to 6 carbon atoms, or a branched chain alkyl of 3 to 8 carbon atoms; and

R<sup>2</sup> is phenyl, naphthyl, anthracyl, phenanthryl, pyridyl, pyrimidyl, triazinyl, furyl, pyrrolyl, pyrazolyl, indolyl, imidazolyl, benzofuryl, benzothienyl, oxazolyl, or thiazolyl each optionally substituted with 0 to 3 substituents selected from straight-chain alkyl of 1 to 6 carbon atoms, branched-chain alkyl of 3 to 8 carbon atoms, alkoxy of 1 to 6 carbon atoms, mono- or dialkylamino of 1 to 6 carbon atoms, nitro, halo, amino, cyano, trifluoromethyl, trifluoromethoxy and hydroxy;

or a pharmaceutically acceptable salt thereof.

Claims 27 to 32 (cancelled)

DOCKET NO.: AM100299 CON US/WYNC-0677

**Application No.:** 10/663,533

Office Action Dated: April 14, 2004

33. (previously presented) A method according to claim 26, wherein said subject is a human.

**PATENT** 

- 34. (previously presented) A method according to claim 26, wherein R<sup>1</sup> is a straight-chained alkyl of 1 to 3 carbon atoms, or a branched chain alkyl of 3 to 6 carbon atoms.
- 35. (previously presented) A method according to claim 26, wherein R<sup>1</sup> is a straight-chained alkyl of 1 or 2 carbon atoms.
- 36. (previously presented) A method according to claim 26, wherein R<sup>2</sup> is phenyl, naphthyl, pyridyl, pyrimidyl, furyl, pyrrolyl, pyrazolyl, indolyl, imidazolyl, benzofuryl, or benzothienyl; each optionally substituted with 1 to 3 substituents the same or different selected from straight-chain alkyl of 1 to 3 carbon atoms, branched-chain alkyl of 3 to 6 carbon atoms, alkoxy of 1 to 3 carbon atoms, mono- or dialkylamino in which each alkyl group has 1 to 3 carbon atoms, nitro, amino, cyano, halogen, trifluoromethyl, trifluoromethoxy, and hydroxy.
- 37. (previously presented) A method according to claim 26, wherein R<sup>2</sup> is phenyl, naphthyl, pyridyl, pyrrolyl, indolyl, or benzothienyl; each optionally substituted with 1 to 3 substituents the same or different selected from nitro, amino, cyano, halogen, trifluoromethyl, trifluoromethoxy, and hydroxy.
- 38. (previously presented) A method according to claim 26, wherein R<sup>2</sup> is trifluoromethylphenyl or methoxyphenyl.
- 39. (previously presented) A method according to claim 26, wherein the R<sup>1</sup>O substituent is bonded to the 1,4-benzodioxan nucleus at the 8 position.

DOCKET NO.: AM100299 CON US/WYNC-0677 PATENT

**Application No.:** 10/663,533

Office Action Dated: April 14, 2004

40. (previously presented) A method according to claim 26, wherein R<sup>1</sup> is a straight-chained alkyl of 1 to 3 carbon atoms, or a branched chain alkyl of 3 to 6 carbon atoms and R<sup>2</sup> is phenyl, naphthyl, pyridyl, pyrimidyl, furyl, pyrrolyl, pyrazolyl, indolyl, imidazolyl, benzofuryl, or benzothienyl; each optionally substituted with 0 to 3 substituents selected from straight-chain alkyl of 1 to 3 carbon atoms, branched-chain alkyl of 3 to 6 carbon atoms, alkoxy of 1 to 3 carbon atoms, mono- or di-alkylamino in which each alkyl group has 1 to 3 carbon atoms, halogen, trifluoromethyl, trifluoromethoxy, and hydroxy.

- 41. (previously presented) A method according to claim 26, wherein R<sup>1</sup> is a straight-chained alkyl of 1 or 2 carbon atoms, and R<sup>2</sup> is phenyl, naphthyl, pyridyl, pyrrolyl, indolyl, or benzothienyl; each optionally substituted with a 0 to 3 substituents selected from nitro, amino, cyano, halogen, trifluoromethyl, trifluoromethoxy, and hydroxy.
- 42. (previously presented) A method according to claim 26, wherein R<sup>1</sup> is a straight chain alkyl of 1 or 2 carbon atoms and R<sup>2</sup> is trifluoromethylphenyl or methoxyphenyl.
- 43. (previously presented) A method according to claim 26, wherein said compound is (S)-8-(8-ethoxy-2,3-dihydrobenzo-[1,4]dioxin-2-ylmethyl)-3-naphthalen-2-yl-8-aza-bicyclo[3.2.1] octan-3-ol or a pharmaceutically acceptable salt thereof.
- 43. (previously presented) A method according to claim 26, wherein said compound is (S)-8-(8-ethoxy-2,3-dihydro-benzo[1,4]dioxin-2-ylmethyl)-3-phenyl-8-aza-bicyclo[3.2.1]octan-3-ol or a pharmaceutically acceptable salt thereof.
- 44. (previously presented) A method according to claim 26, wherein said compound is (S)-3-benzo[b]thiophen-3-yl-8-(8-ethoxy-2,3-dihydro-benzo[1,4]dioxin-2-ylmethyl)-8-aza-bicyclo[3.2.1]octan-3-ol or a pharmaceutically acceptable salt thereof.

DOCKET NO.: AM100299 CON US/WYNC-0677

**Application No.:** 10/663,533

Office Action Dated: April 14, 2004

45. (previously presented) A method according to claim 26, wherein said compound is 8-{[(2S)-8-ethoxy-2,3-dihydrobenzo-[1,4]dioxin-2-yl]methyl)-3-pyridin-2-yl-8-azabicyclo [3.2.1]octan-3-ol or a pharmaceutically acceptable salt thereof.

**PATENT** 

- 46. *(previously presented)* A method according to claim 26, wherein said compound is 8-{[(2S)-8-ethoxy-2,3-dihydrobenzo-[1,4]dioxin-2-yl]methyl)-3-(3-trifluoromethyl-phenyl)-8-aza-bicyclo[3.2.1]octan-3-ol or a pharmaceutically acceptable salt thereof.
- 47. *(previously presented)* A method according to claim 26, wherein said compound is 8-{[(2S)-8-methoxy-2,3-dihydro-1,4-benzodioxin-2-yl]methyl}-3-(2-methoxyphenyl)-8-azabicyclo[3.2.1]octan-3-ol or a pharmaceutically acceptable salt thereof.
- 48. (previously presented) A method according to claim 26, wherein said compound is 8{[(2S)-8-methoxy-2,3-dihydro-1,4-benzodioxin-2-yl]methyl}-3-[3(trifluoromethyl)phenyl]-8-azabicyclo[3.2.1]octan-3-ol or a pharmaceutically acceptable salt thereof.
- 49. *(previously presented)* A method according to claim 26, wherein said compound is 8-{[(2S)-8-methoxy-2,3-dihydro-1,4-benzodioxin-2-yl]methyl}-3-(2-pyridinyl)-8-azabicyclo[3.2.1]octan-3-ol or a pharmaceutically acceptable salt thereof.
- 50. (previously presented) A method according to claim 26, wherein said compound is 3-(1-benzothien-3-yl)-8-{[(2S)-8-methoxy-2,3-dihydro-1,4-benzodioxin-2-yl]methyl}-8-azabicyclo[3.2.1]octan-3-ol or a pharmaceutically acceptable salt thereof.
- 51. (previously presented) A method according to claim 26, wherein said compound is 8-{[(2S)-8-methoxy-2,3-dihydro-1,4-benzodioxin-2-yl]methyl}-3-phenyl-8-azabicyclo[3.2.1]octan-3-ol or a pharmaceutically acceptable salt thereof.

**DOCKET NO.:** AM100299 CON US/WYNC-0677 **PATENT** 

**Application No.:** 10/663,533

Office Action Dated: April 14, 2004

52. (previously presented) A method according to claim 26, wherein said compound is 3- ((2S)-8-methoxy-2,3-dihydrobenzo- [1,4]dioxin-2-ylmethyl)-8-naphthalen-2-yl-3-aza-bicyclo[3.2.1]octan-8-ol or a pharmaceutically acceptable salt thereof.